



# Ball Green Primary School

## Infection Control Policy 2021: Appendix b

### Infection Absence Periods

This table details the minimum required period for staff and pupils to stay away from school following an infection, as recommended by PHE.

\*Identifies a notifiable disease. It is a statutory requirement that doctors report these diseases to their local PHE centre.

Infection	Recommended minimum period to stay away from school	Comments
Athlete's foot	None	Treatment is recommended; however, this is not a serious condition.
Chicken pox	Until all vesicles have crusted over	Follow procedures for vulnerable children and pregnant staff.
Cold sores	None	Avoid contact with the sores.
Conjunctivitis	None	If an outbreak occurs, consult the HPT. In the case of an unmanageable outbreak, the HPT may advise exclusions.
Coronavirus (COVID-19)	Until fully recovered and no other member of the same household is presenting symptoms (10 days if living alone, 14 days if living with others)	If coronavirus is suspected, consult the local HPT.
Diarrhoea and/or vomiting	Whilst symptomatic and 48 hours from the last episode	GPs should be contacted if diarrhoea or vomiting occur after taking part in water-based activities.
Diphtheria*	Exclusion is essential.	Family contacts must be excluded until cleared by the HPT and the HPT must always be consulted.

Infection	Recommended minimum period to stay away from school	Comments
Flu (influenza)	Until recovered	Report outbreaks to the HPT.
Glandular fever	None	
Hand foot and mouth	None	Contact the HPT if a large number of children are affected. Exclusion may be considered in some circumstances.
Head lice	None	Treatment recommended only when live lice seen. Exclusion is not normally permitted. In severe, ongoing cases, the LA does have the power to exclude; however, exclusion should not be overused.
Hepatitis A*	Seven days after onset of jaundice or other symptoms	If it is an outbreak, the HPT will advise on control measures.
Hepatitis B*, C* and HIV	None	Not infectious through casual contact. Procedures for bodily fluid spills must be followed.
Impetigo	48 hours after commencing antibiotic treatment, or when lesions are crusted and healed	Antibiotic treatment is recommended to speed healing and reduce the infectious period.
Measles*	Four days from onset of rash	Preventable by vaccination (MMR). Follow procedures for vulnerable children and pregnant staff.
Meningococcal meningitis*/ septicaemia*	Until recovered	Meningitis ACWY and B are preventable by vaccination.  The HPT will advise on any action needed.
Meningitis* due to other bacteria	Until recovered	Hib and pneumococcal meningitis are preventable by vaccination. The HPT will advise on any action needed.

<b>Infection</b>	<b>Recommended minimum period to stay away from school</b>	<b>Comments</b>
Meningitis viral*	None	As this is a milder form of meningitis, there is no reason to exclude those who have been in close contact with infected persons.
MRSA	None	Good hygiene, in particular environmental cleaning and handwashing, is important to minimise the spread. The local HPT should be consulted.
Mumps*	Five days after onset of swelling	Preventable by vaccination with two doses of MMR.
Ringworm	Exclusion is not usually required	Treatment is required.
Rubella (German measles)	Four days from onset of rash	Preventable by two doses of immunisation (MMR). Follow procedures for pregnant staff.
Scarlet fever	24 hours after commencing antibiotic treatment	Antibiotic treatment is recommended, as a person is infectious for two to three weeks if antibiotics are not administered. If two or more cases occur, the HPT should be contacted.
Scabies	Can return to school after first treatment	The infected person's household and those who have been in close contact will also require treatment.
Slapped cheek/Fifth disease/Parvo Virus B19	None (once rash has developed)	Follow procedures for vulnerable children and pregnant staff.
Threadworms	None	Treatment recommended for the infected person and household contacts.
Tonsillitis	None	There are many causes, but most causes are virus-based and do not require antibiotics.
Tuberculosis (TB)	Pupils with infectious TB can return to school after two weeks of treatment if well enough to do so, and as long as they have responded to anti-TB therapy.	Only pulmonary (lung) TB is infectious. It requires prolonged close contact to spread. Cases with non-pulmonary TB, and cases with pulmonary TB who have effectively completed two weeks of treatment as confirmed by TB nurses, should not be excluded. Consult the local HPT before disseminating information to staff and parents.
Warts and verrucae	None	Verrucae should be covered in swimming pools, gymnasiums and changing rooms.

<b>Infection</b>	<b>Recommended minimum period to stay away from school</b>	<b>Comments</b>
Whooping cough (pertussis)*	Two days from commencing antibiotic treatment, or 21 days from the onset of illness if no antibiotic treatment is given	Preventable by vaccination. Non-infectious coughing can continue for many weeks after treatment. The HPT will organise any necessary contact tracing.

